

The Decoherence of Phase Space Histories

Todd A. Brun

Department of Physics, Caltech, Pasadena, CA 91125

(February 7, 2008)

Abstract

In choosing a family of histories for a system, it is often convenient to choose a succession of locations in phase space, rather than configuration space, for comparison to classical histories. Although there are no good projections onto phase space, several approximate projections have been used in the past; three of these are examined in this paper. Expressions are derived for the probabilities of histories containing arbitrary numbers of projections into phase space, and the conditions for the decoherence of these histories are studied.

I. INTRODUCTION

A great deal of work has been done recently on the use of the decoherence formalism to describe quantum mechanical systems [1–8]. These systems can be described in terms of *decoherent histories*, which can be assigned probabilities obeying classical probability sum rules. While, in principle, a history could be described in terms of any set of variables, most of the work has focussed on histories of particles in configuration space. The simplest and most fine-grained such history is just the classical trajectory of a particle, specifying its exact position at every moment in time. Such histories do not decohere, however. Instead, one must consider considerably coarse-grained histories, in which a position is given only at certain discrete times, and only within certain finite intervals. A history can then be specified by a string of indices α_i , stating which interval the particle is in at time t_i .

Another important class of histories, though, would be descriptions of a system as being in cells of phase space at successive points in time. A small amount of work has been done on this subject [9–13], but they have not been tackled in full generality.

A. The Decoherence Functional

The decoherence functional is a functional on *pairs of histories* of a quantum mechanical system. One simple description of the functional (though not the most general) has the form:

$$D[\alpha, \alpha'] = \text{Tr} \left\{ P_{\alpha_n}^n(t_n) \cdots P_{\alpha_1}^1(t_1) \rho P_{\alpha'_1}^1(t_1) \cdots P_{\alpha'_n}^n(t_n) \right\}. \quad (1.1)$$

In this expression, ρ is the initial density matrix of the system. The $P_{\alpha_i}^i(t_i)$ are Heisenberg *projection operators* onto Hilbert space. At each time t_i these projection operators represent different alternatives for the system. In terms of Schrödinger projections P , these time dependent projections can be written $P(t) = e^{iHt/\hbar} P e^{-iHt/\hbar}$. A complete set of histories has an exhaustive set of alternatives at each time,

$$\sum_{\alpha_i} P_{\alpha_i}^i(t_i) = 1. \quad (1.2)$$

A particular choice of the $\{\alpha_i\}$ represents one particular history, which we will denote α for brevity. Thus, saying that a given history α occurs implies that alternative α_1 occurs at time t_1 , α_2 at time t_2 , and so forth.

The probability of a given history occuring is just given by the diagonal elements of $D[\alpha, \alpha']$:

$$p(\alpha) = D[\alpha, \alpha]. \quad (1.3)$$

In order for these histories to obey the classical probability sum rules, we must require that the set of histories *decoheres*. The usual requirement for this is that the off-diagonal terms of the decoherence function vanish,

$$D[\alpha, \alpha'] = 0, \quad \alpha \neq \alpha'. \quad (1.4)$$

This is actually a sufficient, but not a necessary condition for decoherence. All that is truly required is that the *real parts* of the off-diagonal terms vanish. Most physically decoherent systems, however, display this stronger form of decoherence; we will see this in the cases that we consider.

B. The Transition Matrix

The most common type of problem treated at present is that in which the variables are divided into a *system* and a *reservoir*, or *environment*. In this case one traces over the reservoir variables, and is left with a reduced density matrix on only the system variables. Instead of the simple time evolution operator $e^{-iHt/\hbar}$, the system evolves according to a somewhat more complicated *transition matrix* or *propagator* $\mathbf{T}$. In terms of path integrals this is

$$T(x_f, x'_f, t_f; x_i, x'_i, t_i) = \int \delta x \delta x' \exp \frac{i}{\hbar} \left\{ S[x(t)] - S[x'(t)] + W[x(t), x'(t)] \right\}, \quad (1.5)$$

where the integral is over all paths $x(t)$ and $x'(t)$ from t_i to t_f which begin at x_i and x'_i and end at x_f and x'_f , respectively. $S[x(t)]$ is the action of the system variables independent of the reservoir, and $W[x(t), x'(t)]$ is the Feynman-Vernon *influence phase* arising due to the interactions with the reservoir [14]. The model most commonly considered is one that was developed in the study of Brownian motion [15], in which a one-dimensional particle described by a single variable x interacts with an infinite bath of harmonic oscillators via a linear or weakly nonlinear potential, but this formalism is quite general.

The reduced density matrix $\rho(x; x')$ evolves straightforwardly:

$$\rho(x; x')|_{t_f} = \int dx_i dx'_i T(x_f, x'_f, t_f; x_i, x'_i, t_i) \rho(x_i; x'_i)|_{t_i}. \quad (1.6)$$

Thus, the decoherence functional can now be written

$$D[\alpha, \alpha'] = Tr_x \left\{ P_{\alpha_n}^n \mathbf{T}(\cdots \mathbf{T}(P_{\alpha_1}^1 \rho P_{\alpha'_1}^1) \cdots) P_{\alpha'_n}^n \right\}. \quad (1.7)$$

With the projections P being onto intervals of coordinate space, it is very easy to write the decoherence functional as a constrained path integral over x and x' . For phase-space projections, the form of the decoherence function is more complicated, as we shall see. While there are no true projectors onto cells of phase space (as there are for intervals of coordinate space), there are a number of approximate projectors, and we shall consider these one at a time.

In dealing with phase space, it is natural to consider other representations of the density matrix, most obviously the Wigner distribution:

$$w(X, p) = \frac{1}{\pi} \int d\xi e^{i\xi p/\hbar} \rho(X + \xi/2; X - \xi/2). \quad (1.8)$$

This distribution acts in many ways like a probability distribution in phase space, with the major exception that it can be negative in localized regions. The time evolution of $w(X, p)$ is also described by a transition matrix:

$$w(X_f, p_f)|_{t_f} = \int \int dX_i dp_i T_w(X_f, p_f, t_f; X_i, p_i, t_i) w(X_i, p_i)|_{t_i}. \quad (1.9)$$

$$T_w(X_f, p_f, t_f; X_i, p_i, t_i) = \int \int d\xi_f d\xi_i e^{i(p_f \xi_f - p_i \xi_i)/\hbar} \times T(X_f + \frac{\xi_f}{2}, X_f - \frac{\xi_f}{2}, t_f; X_i + \frac{\xi_i}{2}, X_i - \frac{\xi_i}{2}, t_i), \quad (1.10)$$

where this T is the same transition matrix defined above (1.5). We shall see that the expressions for the probabilities of phase space histories are described very naturally in terms of Wigner distributions.

II. PROBABILITIES OF PHASE SPACE HISTORIES

As mentioned above, there are no true projections onto cells of phase space [10]. This is essentially a side-effect of the uncertainty principle, which prevents both x and p from being localized simultaneously. However, for cells larger in area than $\hbar$, we can approximate projections reasonably well.

In making these calculations, we will find that it is useful to work in terms of $w(X, P)$ and $\mathbf{T}_w$. For this we use the inverses of (1.8) and (1.10), namely

$$\rho(x; x') = \int dp e^{-ip(x-x')/\hbar} w((x+x')/2, p) \quad (2.1)$$

and

$$T(x_f, x'_f, t_f; x_i, x'_i, t_i) = \frac{1}{\pi} \int \int dp_i dp_f e^{ip_i(x_i - x'_i) - ip_f(x_f - x'_f)} \times T_w((x_f + x'_f)/2, p_f, t_f; (x_i + x'_i)/2, p_i, t_i). \quad (2.2)$$

A. Consecutive X and P Projections

While there are no good projections onto phase space cells, projections onto intervals in coordinate or momentum space are perfectly well-defined and straightforward. They are just

$$P_{x_i} = \int_{x_i - \Delta x/2}^{x_i + \Delta x/2} dx |x\rangle\langle x|, \quad (2.3a)$$

$$P_{p_i} = \int_{p_i - \Delta p/2}^{p_i + \Delta p/2} dp |p\rangle\langle p|. \quad (2.3b)$$

We can imagine using a projection to determine which interval of x the system is in, followed a short time later by a projection onto an interval of p . If we let the time between these two “measurements” go to zero, we can make use of the relation

$$\langle x|p\rangle = e^{ipx/\hbar}. \quad (2.4)$$

This is unsatisfactory in a number of ways. The most obvious is that x and p are non-commuting variables, so that P_x and P_p are also non-commuting. The order in which one makes these measurements matters, particularly if the intervals are fairly small (compared to $\hbar$). If we are interested in fairly large cells in phase space, this is of less importance; for classical and quasiclassical systems this is often the case.

Measurements of this type were treated by Halliwell [10]. He also considered another type of two-projection measurement: a pair of successive position measurements, separated by a small time interval Δt , with the momentum determined by the *time of flight* between x_1 and x_2 . I have not considered this type of measurement, as it is ill-defined as $\Delta t \rightarrow 0$, and therefore requires non-trivial consideration of the system’s time evolution between the two position projections. For a system with complex dynamics this is difficult.

The exact projections (2.3) used above are less convenient for the purposes of calculation, though they are certainly more correct. For ease of computation, therefore, it is customary to use approximate Gaussian projections,

$$P_{\bar{x}_i} = \frac{1}{\sqrt{\pi}\Delta x} \int_{-\infty}^{\infty} dx \, e^{-(x-\bar{x}_i)^2/\Delta x^2} |x\rangle\langle x|, \quad (2.5a)$$

$$P_{\bar{p}_i} = \frac{1}{\sqrt{\pi}\Delta p} \int_{-\infty}^{\infty} dp \, e^{-(p-\bar{p}_i)^2/\Delta p^2} |p\rangle\langle p|. \quad (2.5b)$$

Using these projections, the probabilities just reduce to a product of Gaussian integrals, which can, with a little algebra, be easily solved. Using the expressions (2.1) and (2.2) for a history with N measurements of cells of phase space centered on $(\bar{x}_1, \bar{p}_1), (\bar{x}_2, \bar{p}_2), \dots (\bar{x}_N, \bar{p}_N)$, we get

$$P_\alpha = \frac{1}{(\pi\Delta x\Delta p)^{2N}} \int d\{X_i\} d\{p_i\} d\{z_i\} d\{k_i\} d\{z'_i\} d\{k'_i\} d\{\xi_i\}$$

$$\begin{aligned}
& \times W(X_0, p_0) \delta(z_1 - (X_0 + \xi_0/2)) \delta(z'_1 - (X_0 - \xi_0/2)) \\
& \times \exp \left[-\frac{(z_1 - \bar{x}_1)^2 + (z'_1 - \bar{x}_1)^2}{\Delta x^2} - \frac{(k_1 - \bar{p}_1)^2 + (k'_1 - \bar{p}_1)^2}{\Delta p^2} \right. \\
& \left. + \frac{i}{\hbar} \left((k'_1 z'_1 - k_1 z_1) + X_1(k_1 - k'_1) + \xi_1(p_1 - k_1/2 - k'_1/2) - \xi_0 p_0 \right) \right] \\
& \times T(X_2, p_2, t_2; X_1, p_1, t_1) \\
& \times \dots
\end{aligned} \tag{2.6}$$

The integrals over z , z' , k , k' , and ξ are all simple, and yield

$$\begin{aligned}
P_\alpha = & \frac{1}{(2\pi)^{N-1}} \int d\{X_i\} d\{p_i\} w(X_0, p_0) \\
& \times \exp \left[-\frac{2}{\Delta p^2} (p_1 - \bar{p}_1)^2 - \frac{2}{\Delta x^2} (X_0 - \bar{x}_1)^2 - \frac{\Delta x^2}{2\hbar^2} (p_0 - p_1)^2 - \frac{\Delta p^2}{2\hbar^2} (X_0 - X_1)^2 \right] \\
& \times T(X_2, p_2, t_2; X_1, p_1, t_1) \\
& \times \exp \left[-\frac{2}{\Delta p^2} (p_3 - \bar{p}_2)^2 - \frac{2}{\Delta x^2} (X_2 - \bar{x}_2)^2 - \frac{\Delta x^2}{2\hbar^2} (p_2 - p_3)^2 - \frac{\Delta p^2}{2\hbar^2} (X_2 - X_3)^2 \right] \\
& \times T(X_4, p_4, t_3; X_3, p_3, t_2) \\
& \times \dots \\
& \times \exp \left[-\frac{2}{\Delta x^2} (X_{2N-2} - \bar{x}_N)^2 - \frac{\Delta x^2 (p_{2N-2} - \bar{p}_N)^2}{2[\hbar^2 + \Delta x^2 \Delta p^2/4]} \right].
\end{aligned} \tag{2.7}$$

Note that the expression for the probability behaves very reasonably, i.e., the evolution after a “measurement” continues to be centered about the measured values of X and p , with a spread determined by the size of the phase space cell.

A measurement of p followed by a measurement of X produces an expression very similar to the above, and is readily evaluated by the same methods. The differences are chiefly notable when the cell size is small compared to $\hbar$.

B. Coherent State Projections

The closest thing to a true projection onto a cell in phase space is probably the *coherent state* projection $|\bar{p}, \bar{x}\rangle \langle \bar{p}, \bar{x}|$ centered on $(\bar{p}, \bar{x})$. While these are true projections, they are not orthogonal:

$$|\langle \bar{p}, \bar{x} | \bar{p}', \bar{x}' \rangle|^2 = \exp[-(\bar{x} - \bar{x}')^2/2\sigma - \sigma(\bar{p} - \bar{p}')^2/2\hbar^2]. \quad (2.8)$$

Also, these states are overcomplete. Thus, phase space histories built from coherent states cannot be truly decoherent, and can only be even approximately decoherent if a discrete sample of them (e.g., the states corresponding to a lattice of points in phase space) is taken. In a coordinate basis we can represent a coherent state as

$$\langle x | \bar{p}, \bar{x} \rangle = \frac{1}{(\pi\sigma)^{1/4}} \exp\left[-\frac{(x - \bar{x})^2}{2\sigma} + \frac{i\bar{p}x}{\hbar}\right]. \quad (2.9)$$

This expression is useful in evaluating the probability of a coherent state history.

When we consider a history of N “measurements” in phase space using coherent state projections we get an expression analagous to (2.7), which can (again) be solved for the probability:

$$\begin{aligned} P_\alpha = & \frac{4^N}{2\pi\sigma} \int d\{X_i\} d\{p_i\} w(X_0, p_0) \\ & \times \exp\left[-\frac{(X_0 - \bar{x}_1)^2}{\sigma} - \frac{(X_1 - \bar{x}_1)^2}{\sigma} - \frac{(p_0 - \bar{p}_1)^2\sigma}{\hbar^2} - \frac{(p_1 - \bar{p}_1)^2\sigma}{\hbar^2}\right] \\ & \times T(X_2, p_2, t_2; X_1, p_1, t_1) \\ & \times \dots \\ & \times T(X_{2N-2}, p_{2N-2}, t_N; X_{2N-3}, p_{2N-3}, t_{N-1}) \\ & \times \exp\left[-\frac{(X_{2N-2} - \bar{x}_N)^2}{\sigma} - \frac{(p_{2N-2} - \bar{p}_N)^2\sigma}{\hbar^2}\right]. \end{aligned} \quad (2.10)$$

The general behavior of the probabilities is very similar to that in the first case we considered, but even cleaner and easier to see. Coherent states are an excellent way of representing phase space histories.

There is one other kind of approximate projection that we could consider. It is not, in my opinion, a very attractive one, but it has been used in the literature, and so might as well be treated here. Consider approximate “projections” of the form

$$P_{(\bar{p}, \bar{x})} = \frac{1}{\pi\Delta p\Delta x} \int dp dx |p, x\rangle e^{-(p-\bar{p})^2/\Delta p^2 - (x-\bar{x})^2/\Delta x^2} \langle p, x|. \quad (2.11)$$

If we consider N measurements of this form, the probability becomes

$$\begin{aligned}
P_\alpha = & \frac{4\pi\sigma\Delta p\Delta x}{2\sigma + \sigma^2\Delta p^2 + \Delta x^2 + \sigma\Delta p^2\Delta x^2} \int d\{X_i\}d\{p_i\} w(X_0, p_0) \\
& \times \exp\left[-\frac{(X_0 - \bar{x}_1)^2 + (X_1 - \bar{x}_1)^2}{\sigma + \Delta x^2} - \sigma\frac{(p_0 - \bar{p}_1)^2 + (p_1 - \bar{p}_1)^2}{\hbar^2 + \sigma\Delta p^2}\right. \\
& \left. - (\sigma^2\Delta p^2/\hbar^2 + \Delta x^2 + \sigma\Delta p^2\Delta x^2/\hbar^2)\left(\frac{(X_0 - X_1)^2}{2\sigma(\sigma + \Delta x^2)} + \frac{(p_0 - p_1)^2}{2(\hbar^2 + \sigma\Delta p^2)}\right)\right] \\
& \times T(X_2, p_2, t_2; X_1, p_1, t_1) \\
& \times \dots \\
& \times T(X_{2N-2}, p_{2N-2}, t_N; X_{2N-3}, p_{2N-3}, t_{N-1}) \\
& \times \exp\left[-2\frac{\sigma(\sigma + \Delta x^2)(p_{2N-2} - \bar{p}_N)^2 + (\hbar^2 + \sigma\Delta p^2)(X_{2N-2} - \bar{x}_N)^2}{2\sigma\hbar^2 + \sigma^2\Delta p^2 + \Delta x^2\hbar^2 + \sigma\Delta p^2\Delta x^2}\right]. \tag{2.12}
\end{aligned}$$

Again, the same sort of qualitative behavior, but a much uglier expression.

III. DECOHERENCE OF PHASE SPACE HISTORIES

While the above expressions are highly intuitive in their qualitative behavior, we have (in a sense) been putting the cart before the horse. It is meaningless to assign a probability to a history without first being assured that the set of histories described is decoherent. There is nothing in the expressions above to prevent one from choosing extremely tiny cells in phase space, with areas small compared to $\hbar$; yet such histories are certainly not decoherent, as they flagrantly violate the uncertainty principle.

Unfortunately, while we can write *expressions* for the probabilities without having to know much about the physics of the system (i.e., the actual behavior of the transition matrix $\mathbf{T}$), in order to actually calculate them, or to say much about decoherence, we need to know something about the path integrals.

Except in the case of quadratic systems, these integrals are not exactly solvable. Limited treatments of this case have been considered elsewhere [13,16]. Most interesting systems, however, include nonlinearities. This can be handled in one of three ways: numerically; in perturbation theory; or in the semiclassical limit, where solutions are peaked about the “classical trajectory.” The first approach is robust, but does not lend itself to general arguments.

While almost any approach will eventually have to be treated numerically to calculate actual values for probabilities or decoherence functional elements, one would hope to get a rough idea as to a systems behavior before invoking that numerical machinery. The second approach, perturbation theory, is the most commonly adopted. When the nonlinearities are weak, the path integrals can be approximated with considerable precision. Unfortunately, many interesting cases (e.g., chaotic systems) cannot be treated in this fashion; for them, their nonlinearities are intrinsically important. The last approach is limited to systems with sufficient mass and inertia to resist quantum fluctuations [3]. This is useful in considering either the classical limit of quantum systems, or in estimating quantum effects in otherwise classical systems, and is the approach we will adopt here.

Earlier work has concentrated on distinguished systems interacting with a large reservoir or environment whose degrees of freedom can be neglected. As has been shown, such systems give rise to decoherence functionals with probabilities peaked about classical trajectories. The transition matrix for such a system has the form (in the limit of a large thermal reservoir)

$$T(X_1, \xi_1, t_1; X_0, \xi_0, t_0) = \int \delta X \delta \xi \exp \frac{i}{\hbar} \left\{ - \int_{t_0}^{t_1} \left(M \ddot{X}(t) + dV/dX(X(t)) + 2M\gamma \dot{X}(t) - g(t) \right) \xi(t) dt + \frac{iM\gamma kT}{\hbar} \int_{t_0}^{t_1} \xi^2(t) dt + M\xi_1 \dot{X}_1 - M\xi_0 \dot{X}_0 + O(\xi^3) \right\}, \quad (3.1)$$

where X and ξ are variables defined by

$$X = \frac{1}{2}(x + x'), \quad (3.2a)$$

$$\xi = x - x', \quad (3.2b)$$

and the reservoir temperature is T . This is basically a toy system, consisting of a single one-dimensional particle of mass M moving in an arbitrary potential $V(X)$. The interaction with the reservoir provides the dissipative term and a thermal noise; it is this noise which causes the system to decohere. $|\xi|$ is a measure of how far “off-diagonal” the decoherence functional is; for large $|\xi|$ it will clearly be strongly suppressed. This is “medium-strength” decoherence as defined by Gell-Mann and Hartle [2].

Since large ξ is suppressed, we can neglect the higher order terms in ξ with good accuracy. This makes the ξ path integral purely quadratic, and therefore solvable. Doing this integral yields

$$T(X_1, \xi_1, t_1; X_0, \xi_0, t_0) = \sqrt{\frac{\pi \hbar^2}{M \gamma k T}} \int \delta X \exp \left\{ -\frac{1}{M \gamma k T} \int_{t_0}^{t_1} \left(M \ddot{X}(t) + dV/dX(X(t)) + 2M \gamma \dot{X}(t) - g(t) \right)^2 dt + (i/\hbar) M (\xi_1 \dot{X}_1 - \xi_0 \dot{X}_0) \right\}, \quad (3.3)$$

which is clearly peaked about the solution to the classical equation of motion

$$M \ddot{X}(t) + dV/dX(X(t)) + 2M \gamma \dot{X}(t) = g(t), \quad (3.4)$$

more and more strongly in the limit of large M .

Let $X_{\text{cl}}(t)$ be the solution to the above classical equation with the boundary conditions $X_{\text{cl}}(t_0) = X_0$ and $X_{\text{cl}}(t_1) = X_1$. We can then define a new variable $\eta(t)$

$$\eta(t) = X(t) - X_{\text{cl}}(t). \quad (3.5)$$

Clearly $\eta(t)$ has boundary conditions $\eta(t_1) = \eta(t_0) = 0$. As M becomes large, we can treat $\eta(t)$ as a small deviation, and approximate the path integral as

$$T(X_1, \xi_1, t_1; X_0, \xi_0, t_0) = \sqrt{\frac{\pi \hbar^2}{M \gamma k T}} \int \delta \eta \exp \left\{ -\frac{1}{M \gamma k T} \int_{t_0}^{t_1} \left(M \ddot{\eta}(t) + d^2V/dX^2(X_{\text{cl}}(t)) \eta(t) + 2M \gamma \dot{\eta}(t) \right)^2 dt + (i/\hbar) M (\xi_1 \dot{\eta}_1 - \xi_0 \dot{\eta}_0) + (i/\hbar) M (\xi_1 \dot{X}_{\text{cl}}(t_1) - \xi_0 \dot{X}_{\text{cl}}(t_0)) \right\}. \quad (3.6)$$

This path integral is quadratic in η and therefore solvable, at least in principle. This principle runs into a few problems in practice. It assumes that you know $X_{\text{cl}}(t)$ as a function of the boundary conditions. This is true only in very simple cases. In chaotic cases, it may be difficult to determine this function even numerically. Also, this integral contains (in essence) 4th derivatives of η , which complicate the calculation in some ways. Still, by making a few assumptions about the behavior of $X_{\text{cl}}(t)$, we can still extract some useful information from this expression.

Since for the purposes of determining decoherence we are really only interested in the ξ dependence of $\mathbf{T}$, it is straightforward, albeit tedious, to show that

$$T(X_1, \xi_1, t_1; X_0, \xi_0, t_0) = K \exp \left\{ -\frac{M\gamma kT}{\hbar^2} (\lambda_1 \xi_0^2 + \lambda_2 \xi_0 \xi_1 + \lambda_3 \xi_1^2) + (i/\hbar) M (\xi_1 \dot{X}_{\text{cl}}(t_1) - \xi_0 \dot{X}_{\text{cl}}(t_0)) \right\}. \quad (3.7)$$

Note that $\lambda_i = \lambda_i(X_1, X_0, t_1, t_0)$ and $K = K(X_1, X_0, t_1, t_0)$. These functions are not especially easy to calculate, but can be computed numerically if necessary. Simple calculations along those lines seem to show that $\lambda_i/(t_1 - t_0)$ is relatively constant for $(t_1 - t_0)$ short compared to the dynamical time of the system and long compared to the decoherence time, at least for high-probability paths. For longer times, comparable to the dynamic timescale of the system in question, the λ_i vary enormously in magnitude; numerical results showed a variability of more than four orders of magnitude, though most results for $\lambda_1/(t_1 - t_0)$ and $\lambda_3/(t_1 - t_0)$ clustered around certain values, and never became negligibly small. $\lambda_1 \xi_0^2 + \lambda_2 \xi_0 \xi_1 + \lambda_3 \xi_1^2$ is, in any case, always a strictly non-negative quantity. For details of these calculations, see the Appendix.

If we make the (admittedly highly questionable) assumption that the λ_i are roughly constant for constant $(t_1 - t_0)$, then we can estimate the level of decoherence achievable with phase-space projections. For simplicity, we will only look at the projections at a single time:

$$\text{Tr} \left\{ \cdots \mathbf{T}(P_{\alpha_i}^i(t_i) \mathbf{T}(\cdots) P_{\alpha'_i}^i(t_i)) \cdots \right\} \quad (3.8)$$

A. Consecutive X and P Projections

Using the approximate X and P projections described in (2.5b) above, we can examine decoherence by looking at the off-diagonal elements, where the projections are centered on $(\bar{x}, \bar{p})$ and $(\bar{x}', \bar{p}')$ respectively. A single pair of projections at time t_i will multiply the decoherence functional by a factor

$$\int dX_1 dX_2 d\xi_1 d\xi_2 dp dp' T(X_3, \xi_3, t_{i+1}; X_2, \xi_2, t_i)$$

$$\begin{aligned}
& \times \exp \left\{ -\frac{(X_1 + \xi_1/2 - \bar{x})^2}{\Delta x^2} - \frac{(X_1 - \xi_1/2 - \bar{x}')^2}{\Delta x^2} - \frac{(p - \bar{p})^2}{\Delta p^2} - \frac{(p' - \bar{p}')^2}{\Delta p^2} \right. \\
& \left. + i(p - p')(X_2 - X_1) + \frac{i}{2}(p + p')(\xi_2 - \xi_1) \right\} \\
& \times T(X_1, \xi_1, t_i; X_0, \xi_0, t_{i-1})
\end{aligned} \tag{3.9}$$

$$\begin{aligned}
& \sim \int dX_1 dX_2 K^2 \exp \left\{ -\left(\frac{(X_1 - \bar{x})^2 + (X_1 - \bar{x}')^2 + (\Delta p^2 \Delta x^2 / 2\hbar^2)(X_2 - X_1)^2}{\Delta x^2} \right) \right. \\
& + \left(\frac{(8\hbar^2 M\gamma k T \lambda + \hbar^2 \Delta p^2)(\bar{x} - \bar{x}')^2}{2\Delta x^2 (8\hbar^2 M\gamma k T \lambda + \hbar^2 \Delta p^2 + 16(M\gamma k T \lambda)^2 \Delta x^2 + 4M\gamma k T \lambda \Delta p^2 \Delta x^2)} \right) \\
& - \left(\frac{(\hbar^2 + 2M\gamma k T \lambda \Delta x^2)[(p_2 - \bar{p})^2 + (p_2 - \bar{p}')^2]}{8\hbar^2 M\gamma k T \lambda + \hbar^2 \Delta p^2 + 16(M\gamma k T \lambda)^2 \Delta x^2 + 4M\gamma k T \lambda \Delta p^2 \Delta x^2} \right) \\
& - \left(\frac{(2M\gamma k T \lambda \Delta x^2)[(p_1 - \bar{p})^2 + (p_1 - \bar{p}')^2]}{8\hbar^2 M\gamma k T \lambda + \hbar^2 \Delta p^2 + 16(M\gamma k T \lambda)^2 \Delta x^2 + 4M\gamma k T \lambda \Delta p^2 \Delta x^2} \right) \\
& + \left(\frac{(\hbar^2 + 4M\gamma k T \lambda \Delta x^2)(\bar{p} - \bar{p}')^2 - \Delta p^2 (p_2 - p_1)^2}{2(8\hbar^2 M\gamma k T \lambda + \hbar^2 \Delta p^2 + 16(M\gamma k T \lambda)^2 \Delta x^2 + 4M\gamma k T \lambda \Delta p^2 \Delta x^2)} \right) \\
& - i \left(\frac{(4\hbar M\gamma k T \lambda)(\bar{x} - \bar{x}')(\bar{p} + \bar{p}' - 2p_1)}{8\hbar^2 M\gamma k T \lambda + \hbar^2 \Delta p^2 + 16(M\gamma k T \lambda)^2 \Delta x^2 + 4M\gamma k T \lambda \Delta p^2 \Delta x^2} \right) \\
& \left. + \frac{i(\bar{p} - \bar{p}')(X_2 - X_1)}{\hbar} \right\},
\end{aligned} \tag{3.10}$$

where p_1 and p_2 are $M\dot{X}_{cl}$ for boundary conditions $\{X(t_{i-1}) = X_0, X(t_i) = X_1\}$ and $\{X(t_i) = X_2, X(t_{i+1}) = X_3\}$ respectively.

A formidable expression indeed! One can, with difficulty, see that in general this factor will be suppressed for off-diagonal terms. If we simplify matters by taking the semiclassical, high-temperature limit, an examination of the real terms of the exponent show that for $|\bar{x} - \bar{x}'|^2 \sim \delta x^2$ this expression is suppressed by a minimum factor of

$$\exp \left\{ -\frac{\delta x^2}{\Delta x^2} \left(\frac{1}{2} - \frac{\hbar^2}{4M\gamma k T \lambda \Delta x^2} \right) \right\}.$$

A similar examination of the real p terms gives no similar comfort, for we find that there the minimum level of suppression is none at all! This doesn't mean that histories with differing p 's do not decohere; the last imaginary term in the exponent oscillates extremely rapidly, and will tend to suppress all off-diagonal terms as X_1 and X_2 are integrated over. This will work, in general if $|\bar{p} - \bar{p}'|^2 \sim \delta p^2$ is large compared to Δp^2 and $\hbar^2/\Delta x^2$. To

suppress the off-diagonal terms of the decoherence functional by a factor ϵ , where $\epsilon \ll 1$, we must have

$$\begin{aligned}\delta x^2 &\geq -\Delta x^2 \ln \epsilon, \\ \delta p^2 &\geq -\Delta p^2 \ln \epsilon, \\ \Delta x^2 \Delta p^2 &\geq \hbar^2.\end{aligned}\tag{3.11}$$

B. Coherent State Projections

The results from coherent state projections are similar, but somewhat cleaner and easier to see. In this case the factor from the projections at one time t_i goes as

$$\begin{aligned}\int dX_1 dX_2 K^2 \exp \Big\{ & - \left(\frac{(X_1 - \bar{x})^2 + (X_1 - \bar{x}')^2 + (X_2 - \bar{x})^2 + (X_2 - \bar{x}')^2}{2\sigma} \right) \\ & + \left(\frac{\hbar^2(\bar{x} - \bar{x}')^2}{2\hbar^2\sigma + 8\hbar M\gamma kT\lambda\sigma^2} \right) \\ & - \left(\frac{(p_1 - \bar{p})^2 + (p_1 - \bar{p}')^2 + (p_2 - \bar{p})^2 + (p_2 - \bar{p}')^2 - (\bar{p} - \bar{p}')^2}{2\hbar^2/\sigma + 8\hbar M\gamma kT\lambda} \right) \\ & - \frac{i\hbar(\bar{x} - \bar{x}')(p_2 - p_1)}{\hbar^2 + 4M\gamma kT\lambda\sigma} + \frac{i(\bar{p} - \bar{p}')(X_2 - X_1)}{\hbar} \Big\}.\end{aligned}\tag{3.12}$$

Here again, we see that in the semiclassical limit this reduces to a minimal level of suppression

$$\exp \left\{ -\frac{\delta x^2}{\sigma} \left(\frac{1}{2} - \frac{\hbar^2}{8M\gamma kT\lambda\sigma} \right) \right\}$$

for the x terms, and that δp^2 must be large compared to $\Delta p^2 \geq \hbar^2/\sigma$.

IV. CONCLUSIONS

While no set of approximate phase space projections treated in this paper is completely satisfactory, they do serve to illustrate certain traits that phase space histories should possess. Highly discontinuous trajectories are suppressed, and as one goes to the semiclassical

limit the probabilities of histories become peaked about the classical solutions. While precise statements about decoherence are hard to make, given the difficulty of solving the problem for highly general systems, rough arguments indicate that the size of phase space cells needed for decoherence is much larger than that naively indicated by the uncertainty principle ($\Delta x \Delta p \sim \hbar$).

ACKNOWLEDGMENTS

The guidance of Murray Gell-Mann is gratefully acknowledged. I had helpful conversations and exchanges with many people; among the most notable were Seth Lloyd, Jim Hartle, Jonathon Halliwell, Carlton Caves, and Jeff Kimble. Much of the algebra (though not all, alas!) was made easier by use of the Mathematica symbolic mathematics program. This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG03-92-ER40701.

APPENDIX: PATH INTEGRAL FOR THE TRANSITION MATRIX

The path integral in (3.6) is somewhat unusual in that it has only two boundary conditions ($\eta(t_0) = \eta(t_1) = 0$) for an integrand with four derivatives! Thus, the usual prescription for solving quadratic path integrals is not immediately applicable.

This procedure can still be used, however, by the simple expedient of imposing two more boundary conditions, $\dot{\eta}(t_0) = v_0$ and $\dot{\eta}(t_1) = v_1$, and solving the path integral, then integrating the result over all values of v_0 and v_1 .

The path integral to be solved is then

$$\int \delta\eta \exp\left\{S[\eta(t)] + i(M/\hbar)(\xi_1 v_1 - \xi_0 v_0)\right\} = F(t_1, t_0) \exp\{S_{cl} + i(M/\hbar)(\xi_1 v_1 - \xi_0 v_0)\}, \quad (\text{A1})$$

where

$$S[\eta(t)] = -\frac{1}{M\gamma kT} \int_{t_0}^{t_1} \varepsilon^2(t) dt, \quad (\text{A2})$$

$$\varepsilon(t) = M(\ddot{\eta} + 2\gamma\dot{\eta} + f(t)\eta), \quad (\text{A3})$$

$$f(t) = \frac{1}{M} \frac{d^2 V}{dX^2}(X_{\text{cl}}(t)), \quad (\text{A4})$$

and $F(t_1, t_0)$ is an undetermined multiplier independent of the boundary conditions of η . (Of course, since $f(t)$ is defined in terms of $X_{\text{cl}}(t)$, this whole solution *is* dependent on X_0 and X_1 . This dependence is complicated, as we will see.)

The classical action S_{cl} is the action of the path $\eta(t)$ that obeys the classical equation of motion. For

$$S = \int_{t_0}^{t_1} L(\eta, \dot{\eta}, \ddot{\eta}) dt$$

the Euler-Lagrange equation is

$$\frac{d^2}{dt^2} \left(\frac{\partial L}{\partial \ddot{\eta}} \right) - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\eta}} \right) + \left(\frac{\partial L}{\partial \eta} \right) = 0. \quad (\text{A5})$$

Plugging in our definitions for $S[\eta(t)]$ and $\varepsilon(t)$, we get the equations

$$\ddot{\eta} + 2\gamma\dot{\eta} + f(t)\eta = \varepsilon(t)/M, \quad (\text{A6a})$$

$$\ddot{\varepsilon} - 2\gamma\dot{\varepsilon} + f(t)\varepsilon = 0. \quad (\text{A6b})$$

The first of these follows from the definition (A3), the second from (A5).

The solution to the first equation is

$$\varepsilon(t) = M e^{\gamma t} (A a_1(t) + B a_2(t)), \quad (\text{A7})$$

where a_1 and a_2 are two independent solutions of the equation

$$\ddot{a} = (\gamma^2 - f(t))a. \quad (\text{A8})$$

This equation is not easily solved analytically in most cases. For slowly varying $f(t)$ one can approximate the solution

$$a(t) \approx \exp\left\{\pm \int_{t_0}^t \sqrt{\gamma^2 - f(s)} ds\right\}; \quad (\text{A9})$$

in any case, (A8) is readily solvable numerically. While any independent boundary conditions will work for a_1 and a_2 , a convenient choice is

$$\begin{aligned} a_1(0) &= 1, \dot{a}_1(0) = 0; \\ a_2(0) &= 0, \dot{a}_2(0) = 1. \end{aligned} \quad (\text{A10})$$

This is then plugged into the equation for η to give the solution

$$\begin{aligned} \eta(t) &= e^{-\gamma t}(Ca_1(t) + Da_2(t)) \\ &+ (e^{-\gamma t}/M) \int_{t_0}^t \left(\frac{a_1(t)a_2(s) - a_2(t)a_1(s)}{\dot{a}_1(s)a_2(s) - \dot{a}_2(s)a_1(s)} \right) \varepsilon(s) e^{2\gamma s} ds. \end{aligned} \quad (\text{A11})$$

Imposing the boundary conditions on η then gives us a set of equations involving A , B , C , and D :

$$C = 0, \quad (\text{A12a})$$

$$D = v_0, \quad (\text{A12b})$$

$$v_0 a_2(t_1) + A(a_1(t_1)I_2 - a_2(t_1)I_1) + B(a_1(t_1)I_3 - a_2(t_1)I_2) = 0, \quad (\text{A12c})$$

$$v_0 \dot{a}_2(t_1) + A(\dot{a}_1(t_1)I_2 - \dot{a}_2(t_1)I_1) + B(\dot{a}_1(t_1)I_3 - \dot{a}_2(t_1)I_2) = v_1 e^{\gamma t_1}, \quad (\text{A12d})$$

where

$$I_1 = \int_{t_0}^{t_1} \frac{a_1(t)^2 e^{2\gamma t} dt}{\dot{a}_1(t)a_2(t) - \dot{a}_2(t)a_1(t)}, \quad (\text{A13})$$

$$I_2 = \int_{t_0}^{t_1} \frac{a_1(t)a_2(t) e^{2\gamma t} dt}{\dot{a}_1(t)a_2(t) - \dot{a}_2(t)a_1(t)}, \quad (\text{A14})$$

$$I_3 = \int_{t_0}^{t_1} \frac{a_2(t)^2 e^{2\gamma t} dt}{\dot{a}_1(t)a_2(t) - \dot{a}_2(t)a_1(t)}. \quad (\text{A15})$$

Again, these integrals are usually only solvable numerically.

Solving for A and B then yields

$$\begin{aligned} A &= \frac{(\dot{a}_1(t_1)a_2(t_1) - a_1(t_1)\dot{a}_2(t_1))I_3 v_0 + e^{\gamma t_1}(a_1(t_1)I_3 - a_2(t_1)I_2)v_1}{(a_1(t_1)\dot{a}_2(t_1) - \dot{a}_1(t_1)a_2(t_1))(I_2^2 - I_1 I_3)} \\ &= A_0 v_0 + A_1 v_1, \end{aligned} \quad (\text{A16})$$

$$\begin{aligned}
B &= \frac{-(\dot{a}_1(t_1)a_2(t_1) - a_1(t_1)\dot{a}_2(t_1))I_2v_0 - e^{\gamma t_1}(a_1(t_1)I_2 - a_2(t_1)I_1)v_1}{(a_1(t_1)\dot{a}_2(t_1) - \dot{a}_1(t_1)a_2(t_1))(I_2^2 - I_1I_3)} \\
&= B_0v_0 + B_1v_1.
\end{aligned} \tag{A17}$$

Plugging these results back into the definitions of $\varepsilon(t)$ and $S[\eta(t)]$ gives a value for the path integral

$$\begin{aligned}
&\exp\left\{S_{\text{cl}} + (iM/\hbar)(\xi_1v_1 - \xi_0v_0)\right\} = \\
&\exp\left\{-\frac{M}{\gamma kT}(A^2I_4 + 2ABI_5 + B^2I_6) + i(M/\hbar)(\xi_1v_1 - \xi_0v_0)\right\},
\end{aligned} \tag{A18}$$

where

$$I_4 = \int_{t_0}^{t_1} a_1(t)^2 e^{2\gamma t} dt, \tag{A19}$$

$$I_5 = \int_{t_0}^{t_1} a_1(t)a_2(t)e^{2\gamma t} dt, \tag{A20}$$

$$I_6 = \int_{t_0}^{t_1} a_2(t)^2 e^{2\gamma t} dt. \tag{A21}$$

Clearly, the exponent is quadratic in v_0 and v_1 . Integrating over these two boundary conditions, we perform two gaussian integrals, and arrive at our final result

$$\begin{aligned}
&\int dv_0 dv_1 \int \delta\eta \exp\left\{S[\eta(t)] + i(M/\hbar)(\xi_1v_1 - \xi_0v_0)\right\} = \\
&K \exp\left\{-\frac{M\gamma kT}{\hbar^2}(\lambda_1\xi_0^2 + \lambda_2\xi_0\xi_1 + \lambda_3\xi_1^2)\right\}.
\end{aligned} \tag{A22}$$

where

$$\lambda_1 = \frac{A_1^2I_4 + 2A_1B_1I_5 + B_1^2I_6}{4(A_0B_1 - A_1B_0)^2(I_5^2 - I_4I_6)}, \tag{A23a}$$

$$\lambda_2 = \frac{A_0A_1I_4 + (A_1B_0 + A_0B_1)I_5 + B_0B_1I_6}{2(A_0B_1 - A_1B_0)^2(I_5^2 - I_4I_6)}, \tag{A23b}$$

$$\lambda_3 = \frac{A_0^2I_4 + 2A_0B_0I_5 + B_0^2I_6}{4(A_0B_1 - A_1B_0)^2(I_5^2 - I_4I_6)}, \tag{A23c}$$

$$K = \frac{\pi\gamma kT}{M\sqrt{(A_1B_0 - A_0B_1)^2(I_4I_6 - I_5^2)}} F(t_1, t_0). \tag{A23d}$$

Clearly, these quantities are dependent on X_0 and X_1 : $\lambda_i = \lambda_i(X_0, X_1, t_0, t_1)$ and $K = K(X_0, X_1, t_0, t_1)$. In principle, K can be calculated, but in practice it is not necessary to do so in order to make arguments about decoherence. It would be necessary to do so in order to actually compute the probability of a history.

The derivations in this section, tedious as they are, can nevertheless be readily automated. Once one determines the classical trajectory X_{cl} which corresponds to the boundary conditions X_0 and X_1 , determining the λ_i numerically is straightforward. I have used this technique to examine the values of the λ_i for the forced, damped duffing oscillator model. For long times $(t_1 - t_0)$ the λ_i varied enormously in magnitude as a function of X_0 and X_1 . For times short compared to the dynamical time of the system, however, the λ_i were nearly linear: $\lambda/(t_1 - t_0) \approx 0.085$. For some boundary conditions they might become considerably larger, but in what was, admittedly, not an exhaustive sampling, none got much smaller.

REFERENCES

- [1] M. Gell-Mann and J. Hartle, in *Complexity, Entropy, and the Physics of Information*, SFI Studies in the Sciences of Complexity v.8, ed. by W. Zurek, Addison-Wesley, Reading (1990).
- [2] M. Gell-Mann and J. Hartle, Proceedings of the 25th International Conference on High-Energy Physics, Singapore, 1303 (August 1990).
- [3] M. Gell-Mann and J. Hartle, Phys. Rev. D **47**, 3345 (1993).
- [4] R. Omnès, Rev. Mod. Phys. **64**, 339 (1992).
- [5] W. H. Zurek, Phys. Rev. D **24**, 1516 (1981).
- [6] W. H. Zurek, Physics Today **46**, 81 (1993).
- [7] H. F. Dowker and J. J. Halliwell, Phys. Rev. D **46**, 1580 (1992).
- [8] T. Brun, Phys. Rev. D **47**, 3383 (1993).
- [9] T. Brun, Caltech preprint CALT-68-1882 (1993).
- [10] J. J. Halliwell, Phys. Rev. D **46**, 1610 (1992).
- [11] J. J. Halliwell, Phys. Rev. D **48**, 2739 (1993).
- [12] J. J. Halliwell, Phys. Rev. D **48**, 4785 (1993).
- [13] J. Twamley, Phys. Rev. D **48**, 5730 (1993).
- [14] R. P. Feynman and F. L. Vernon, Ann. Phys. (N.Y.) **24**, 118 (1963).
- [15] A. O. Caldeira and A. J. Leggett, Physica **121A**, 587 (1983)
- [16] J. Twamley, Phys. Rev. A **48**, 2627 (1993).